

.....

Safety Assessment: Synthetic Cannabidiol (CBD) as a novel for use in food supplements.

.....

Date 29th April 2024

<https://doi.org/10.46756/sci.fsa.fei457>

Executive Summary

An application was submitted to the Food Standards Agency (FSA) and Food Standards Scotland (FSS) in January 2021 from Chanelle McCoy CBD Ltd (“the applicant”) for the authorisation of synthetic cannabidiol (CBD) as a novel food.

The novel food is a synthetic >98% pure form CBD which is intended to be used as food supplements for adults.

For CBD a provisional Acceptable Daily Intake (ADI) of 10 mg/day has been published by the FSA and was considered in assessing this novel food. The provisional ADI (section 2.7) was recommended, subject to the existing advice to consumers that pregnant and breastfeeding women and people taking any prescription medication should avoid the consumption of CBD. Consumers on regular medications should seek advice from a medical professional before using any type of CBD food product. In addition, children and prospective parents trying for a baby are advised against consumption of CBD, as are those who are immunosuppressed, due to remaining data gaps and residual uncertainties concerning the safety of CBD for these groups of consumers. These contraindications would also apply to this novel food.

To support the FSA and FSS in their evaluations of the application, the Advisory Committee on Novel Foods and Processes (ACNFP) were asked to review the dossier and supplementary information provided by the applicant. Please note the Committee did not consider any potential health benefits or claims arising from consuming the food, as the focus of the novel food assessment is to ensure the food is safe, and not putting consumers at a nutritional disadvantage.

The FSA and FSS concluded based on the advice of the ACNFP, that the applicant had provided sufficient information to assure the novel food, synthetic CBD, was safe under the proposed conditions of use. The anticipated intake levels and the proposed use in foods and food supplements was not considered to be nutritionally disadvantageous.

The views of the ACNFP have been taken into account in the regulatory assessment which represents the opinions of the FSA and FSS.

Contents

Safety Assessment: Synthetic Cannabidiol (CBD) as a novel for use in food supplements.	0
Executive Summary	1
List of tables	4
Abbreviations	4
1. Introduction	6
2. Assessment	7
2.1 Identity of the novel food	7
2.2 Production Process	8
2.3 Compositional information	10
2.4 Stability	18
2.5 Specification	19
2.6 History of Use	21
2.7 Proposed Use and Anticipated Intake	22
2.8 Absorption, Distribution, Metabolism and Excretion (ADME)	24
2.9 Nutritional information	25
2.10 Toxicological information	26
2.11 Allergenicity	28
3. Discussion	28
4. Conclusion	30
Acknowledgements	30
References	32

List of tables

Table 1: Comparison of information on permitted residue levels for solvents used in the novel foods production compared to the proposed specification.

Table 2: Compositional analysis of representative batches of synthetic cannabidiol (CBD)

Table 3: The content of heavy metals, minerals and trace elements in the novel food

Table 4: The microbiological analysis of the novel food (synthetic CBD)

Table 5: Water activity analysis of the novel food (synthetic CBD)

Table 6: Specification of the novel food

Table 7: The proposed uses and maximum use levels for the novel food.

Abbreviations

Acronym	Definition
¹ H NMR	¹ H (proton) nuclear magnetic resonance
ACMD	Advisory Council on the Misuse of Drugs
ACNFP	Advisory Committee on Novel Foods and Processes
ADI	Acceptable Daily Intake
ADME	Absorption, Distribution, Metabolism and Excretion
ARfD	Acute Reference Dose
a _w	Water activity
bw	Body weight
CAS	Chemical Abstracts Service
CBD	Cannabidiol
C _{max}	Peak serum concentration
COT	Committee on Toxicity
CFU	Colony Forming Unit
DAD	Diode Array Detector

Acronym	Definition
EC	European Commission
EFSA	European Food Safety Agency
EMA	Environmental Medicines Agency
EU	European Union
FDA	Food and Drug Administration (USA)
FSA	Food Standards Agency
FSS	Food Standards Scotland
GC	Gas Chromatography
GLP	Good Laboratory Practice
HACCP	Hazards Analysis and Critical Control Points
HPLC	High-performance liquid chromatography
IR	Infra-red
LOAEL	Lowest Observable Adverse Effect Level
NOAEL	No Observable Adverse Effect Level
NM	Not measured
OECD	Organisation for Economic Co-operation and Development
Tmax	Time to maximum concentration
USP	United States Pharmacopeia
UV	Ultra-violet

1. Introduction

1. FSA and FSS have undertaken safety assessments for synthetic CBD under the novel foods legislation, assimilated Regulation (EU) 2015/2283. To support the safety assessment, the ACNFP provided the advice outlined in this opinion to the FSA and FSS.
2. The evaluation by the ACNFP assessed the food safety risks of the novel food and its production, in line with Article 7 of assimilated Commission Implementing Regulation (EU) 2017/2469. The regulatory framework and the technical guidance put in place by the European Food Safety Agency (EFSA) for full novel food applications is used as the basis and structure for the assessment (EFSA NDA Panel, 2021).
3. An application was submitted to the Food Standards Agency (FSA) and Food Standards Scotland (FSS) in January 2021 from Chanelle McCoy Ltd. (“the applicant”) for the authorisation of synthetic cannabidiol (CBD) as a novel food. The novel food is a synthetic >98% pure form CBD which is intended to be used as a food supplement for adults.
4. Advice was sought from the joint Subgroup of the ACNFP and the Committee on Toxicity (COT) on CBD and hemp derived products, on the quality of the toxicological evidence submitted to support the application. The ACNFP and COT have issued a joint statement on a provisional ADI that can be applied to CBD ingredients of >98% purity. This, and wider evidence available in the public domain, was taken into account in reviewing the toxicological evidence for this application.
5. The final advice from the Committee was agreed at the 164th meeting, allowing the FSA and FSS to complete the risk assessment.

6. The document outlines the conclusions of the FSA and FSS on the safety of a synthetic >98% pure form CBD (as detailed in application RP07) as a novel food.

2. Assessment

2.1 Identity of the novel food

7. The novel food is a synthetic >98% pure form of CBD which is a white to slightly beige crystalline powder of experimentally derived $100\% \pm 2\%$ purity. CBD is found in natural form in hemp as the (-)- trans isomer. As this CBD was synthesised chemically, chirality analysis of the compound by a commercially validated chiral HPLC method demonstrated a chiral purity of no less than 99.5% (-) - trans cannabidiol. Information to support stereoselective synthesis, and characterisation, was provided for six batches of the novel food.
8. CBD in this application is characterised by the chemical formula: $C_{21}H_{30}O_2$; molecular mass: 314.46 g/mol; CAS number: 13956-29-1; isomer: (-)-trans-cannabidiol; IUPAC name: 2-[(1R,6R)-3-methyl-6-prop-1-en-2-ylcyclohex-2-en-1-yl]-5-pentylbenzene-1,3-diol.




Diagram 1: The molecular structure of CBD

9. Confirmation of its identity was provided by Infrared (IR) spectroscopy, 1H Nuclear Magnetic Resonance (NMR) spectroscopy, mass spectrometry, and Ultraviolet-visible spectroscopy (UV-Vis).

2.2 Production Process

10. The synthetic >98% pure form of CBD is manufactured using a stereoselective multi-step process under controlled conditions.
11. Certificates of analysis for raw starting materials used in the chemical synthesis were provided to demonstrate the effectiveness of the controls at this point in the process.
12. This process begins with olivetol and menthadienol as the primary starting materials, which undergo mixing, filtration, and distillation steps with catalysts present, resulting in the formation of the desired synthetic >98% pure CBD form.
13. For optional purification steps for commercial purposes, the crude final CBD ingredient is washed and separated several times using various processing aids.
14. The ACNFP considered whether the use of solvents as processing aids left any residues that needed to be flagged to risk managers. Comparison was made to residue limits for other consumed products as detailed in Table 1. Residues of solvents have been included in the specification.

Table 1: Comparison of information on permitted residue levels for solvents used in the novel foods production compared to the proposed specification.

Solvent used	Available data on safe maximum level of consumption	Level in specification for the novel food
Dichloromethane	Guidance on residues in Pharmaceutical products sets a Permissible Daily Exposure of 6.0 mg/day or at a concentration of 600ppm ¹ Permitted extraction solvent residue on coffee at 2 mg/kg and 5 mg/kg in tea ²	600 mg/kg CBD
Methanol	Guidance on residues in pharmaceutical products sets Permissible Daily Exposure of 30.0 mg/day or a concentration of 3000 ppm. ³ Extraction solvent residue maximum level of 10 mg/kg product for all food uses ² .	3000 mg/kg CBD

[1 Q3C \(R8\) Step 5 - impurities: guideline for residual solvents \(europa.eu\)](#)

[2 Directive 2009/32/EC of the European Parliament and of the Council of 23 April 2009 on the approximation of the laws of the Member States on extraction solvents used in the production of foodstuffs and food ingredients](#)

[3 Q3C \(R8\) Step 5 - impurities: guideline for residual solvents \(europa.eu\)](#)

Solvent used	Available data on safe maximum level of consumption	Level in specification for the novel food
Isopropanol	Extraction solvent residue maximum level of 10 mg/kg ² ADI of 2.4 mg/kg bw as acceptable previous cargoes for edible fats and oils ⁴	5000 mg/kg CBD

15. The evidence presented (see Table 2 below) on composition indicates compliance with the specification for residues of solvents. When considered at the level of consumption the evidence suggests the levels of solvent residues in the novel food are below those which would represent a safety concern.
16. A HACCP statement was provided along with further details of the process and how it operates. The production process has characterised the potential hazards and detailed the corresponding control measures sufficiently.

2.3 Compositional information

17. Results from six independent batches of the novel food demonstrated that the CBD content is produced consistently (Table 2).

⁴ Scientific Opinion on the evaluation of the substances currently on the list in the annex to Commission Directive 96/3/EC as acceptable previous cargoes for edible fats and oils – Part II of III

Table 2: Compositional analysis of representative batches of synthetic cannabidiol (CBD)

Parameter	Method of Analysis	Batch 1	Batch 2	Batch 3	Batch 4	Batch 5	Batch 6
Identity and Composition	-----	-----	-----	-----	-----	-----	-----
Appearance	Visual	White crystalline powder	White crystalline powder	Almost white crystalline powder	White to slightly beige crystalline powder	White to slightly beige crystalline powder	White to slightly beige crystalline powder
Identification by IR	USP 197	Corresponds to reference	Corresponds to reference	Corresponds to reference	Corresponds to reference	Corresponds to reference	Corresponds to reference
Identification by HPLC	In house method	Corresponds to reference	Corresponds to reference	Corresponds to reference	Corresponds to reference	Corresponds to reference	Corresponds to reference
Water content (%)	USP 921	≤0.05	≤0.05	≤0.05	0.0	0.2	0.0
Residue on ignition (%)	USP 281	≤0.05	≤0.05	≤0.05	0.1	0.1	0.0

Parameter	Method of Analysis	Batch 1	Batch 2	Batch 3	Batch 4	Batch 5	Batch 6
Identity and Composition	-----	-----	-----	-----	-----	-----	-----
Specific optical rotation (degrees)	USP 781	-128	-130	-132	-129	-129	-129
CBD (%)	HPLC (in house method)	99.4	99.8	99.9	100.5	100.7	100.2
Olivetol content (%)	HPLC (in house method)	≤0.03	≤0.03	≤0.03	≤0.03	≤0.03	≤0.03
4-monobromo-cannabidiol (%)	HPLC (in house method)	≤0.03	≤0.03	≤0.03	≤0.03	≤0.03	≤0.03
Delta-9-tetrahydrocannabidiol (%)	HPLC (in house method)	≤0.02	≤0.02	≤0.02	≤0.02	≤0.02	≤0.02

Parameter	Method of Analysis	Batch 1	Batch 2	Batch 3	Batch 4	Batch 5	Batch 6
Identity and Composition	-----	-----	-----	-----	-----	-----	-----
Individual unspecified impurities (%)	HPLC (in house method)	≤0.05	≤0.05	≤0.05	≤0.05	≤0.05	≤0.05
Total impurities (%)	HPLC (in house method)	≤0.05	≤0.05	≤0.05	≤0.05	≤0.05	≤0.05
Residues (mg/kg of CBD)	-----	-----	-----	-----	-----	-----	-----
Methanol	GC (in house method)	≤300	≤300	≤300	NM	NM	NM
<i>n</i> -heptane	GC (in house method)	≤500	≤500	≤500	≤500	≤500	≤500
Dichloromethane	GC (in house method)	≤60	≤60	≤60	≤60	≤60	≤60

Parameter	Method of Analysis	Batch 1	Batch 2	Batch 3	Batch 4	Batch 5	Batch 6
Residues (mg/kg of CBD)	-----	-----	-----	-----	-----	-----	-----
Triethylamine	GC (in house method)	≤500	≤500	≤500	≤500	≤500	≤500
Isopropanol	GC (in house method)	≤500	≤500	≤500	≤500	≤500	≤500
Isooctane	GC (in house method)	≤500	≤500	≤500	≤500	≤500	545

2 GC = Gas chromatography; HPLC = high-performance liquid chromatography; IR = infrared spectroscopy; NM = not measured; USP

3 = United States Pharmacopeia

4

18. CBD content is consistently above 98% pure with negligible amounts of other cannabinoids detected across the six representative batches.
19. It is recognised that the detection and characterisation of cannabinoids in a range of food matrices is an evolving area and there are yet to be internationally recognised methods. The limitations of analytical methodology available have been subject to discussion in the Joint ACNFP and COT CBD Subgroup and remains a source of uncertainty in the assessment. As a result, the robustness, accuracy, and precision of the methods have been considered in interpreting the data on THC and were considered appropriate in this case.
20. Delta-9-tetrahydrocannabinidiol (THC), as a potential contaminant in the novel food, was detected at levels at or below 0.02 mg/kg CBD in all six batches tested.
21. A literature review was undertaken as part of the assessment of CBD as a novel food, to understand the impact on the safety of foods with trace levels of contamination with THC. The joint ACNFP and COT subgroup reviewed the information from literature and identified a point of departure from the European Food Safety Authority (EFSA) opinion on THC as a contaminant in milk and meat (EFSA, 2015). The point of departure was a LOAEL (Lowest Observed Adverse Effect Level) of 0.036 mg/kg/bw/day from the most sensitive individuals and the lowest dose tested in the human clinical studies reviewed.
22. Uncertainty factors, including a factor of 3 to extrapolate from a LOAEL to a NOAEL (No Observed Adverse Effect Level), were applied to the LOAEL identified. This was considered appropriate as the effects are mild to moderate in severity. A further factor of 10 was applied for person-to-person variation, resulting in a total uncertainty factor of 30. The factor of 30 applied to the 0.036 mg/kg/bw/day LOAEL results in a level of 1 µg /kg bw/day that would be represent a safe upper intake for consumed THC as a contaminant in food. This was identified an acute reference dose (ARfD) (EFSA, 2015).
23. The Subgroup agreed the ARfD to be sufficiently protective to apply to the UK population. It was noted that in applying the acute reference dose EFSA have

assumed that the effects seen would be the same if humans were exposed to multiple doses of THC at very low levels. The Subgroup commented that there was no data to verify this assumption, but if setting limits the dataset is the best available.

24. The levels of THC in the novel food, once adjusted to take account the proposed uses – 10 mg of CBD being consumed a day, were below the ARfD identified by EFSA of 1 µg /kg bw/day or 70 µg/day for a healthy adult. This level does not present a concern in terms of consumer safety for the novel food under the proposed conditions of use.
25. To ensure THC levels remain consistently low in the production of CBD, THC should be a standard substance included in the specification as relevant to all batches produced.
26. Analytical data, presented as the mean of three independent batches of the novel food, demonstrated that heavy metals where detected were present in very low levels and were below established UK regulatory limits where applicable (arsenic, cadmium, mercury and lead) (Table 3).

Table 3: The content of heavy metals, minerals and trace elements in the novel food

Parameter (mg/kg)	Maximum concentration measured (mg/kg)
Arsenic	≤0.1
Cadmium	≤0.1
Mercury	≤0.1
Lead	≤0.1
Cobalt	≤0.1
Nickel	0.2
Vanadium	≤0.1
Silver	≤0.1
Gold	≤0.2
Iridium	≤0.1

Parameter (mg/kg)	Maximum concentration measured (mg/kg)
Osmium	≤0.7
Palladium	≤0.1
Platinum	≤0.1
Rhodium	≤0.1
Ruthenium	≤0.1
Selenium	≤0.1
Thallium	≤0.1
Barium	≤1
Chromium (III)	0.1
Copper	0.1
Lithium	≤0.1
Molybdenum	≤0.4
Antimony	≤0.1
Tin	≤0.1

27. Analytical data concerning the microbiological content from three independent batches of the novel food were reported (Table 4). The process in manufacturing this novel food uses extreme high and low temperatures, harsh pH conditions and alcohol solvents. Full microbial risk assessment as per USP 61 and USP 62 confirm that the novel food does not raise a safety concern and consistently meets the proposed microbial specification levels.

Table 4: The microbiological analysis of the novel food (synthetic CBD)

Parameter	Batch 1	Batch 2	Batch 3
Total aerobic microbial plate count	< 100 CFU/g	< 100 CFU/g	< 100 CFU/g
Total yeast and mould plate count	< 100 CFU/g	< 100 CFU/g	< 100 CFU/g

28. Additionally, water activity of three representative samples of the novel food was measured as per USP 1112. The results further demonstrate that microbial growth in the CBD novel food is unlikely at the low a_w values (below 0.6) for the final ingredient (Table 5). Therefore, microbial analysis is not a required parameter for inclusion in the specification.

Table 5: Water activity analysis of the novel food (synthetic CBD)

Batch	Result (a_w)
1	0.40
2	0.40
3	0.27
Average	0.36

29. The data presented indicated the novel food was appropriately characterised.

2.4 Stability

30. The stability of the synthetic CBD was assessed under real-time conditions (25 °C and 60% relative humidity) and under refrigerated conditions (5 °C) in 3 batches for 3 months, in a further 3 batches for 6 months and another 3 batches for 9 months. Results showed that the novel food meets the specification criteria for CBD content, and no changes in appearance, water content and impurity levels are seen over these time periods. The THC content was also tested using HPLC

and remained consistently below 0.05% w/w across the time period which is below the 0.10% w/w in the specification.

31. The stability of the synthetic CBD was assessed under accelerated conditions (40°C and 75% relative humidity) in six batches over a maximum period of 9 months. Results confirmed that the novel food meets the specification criteria for CBD content and no changes in appearance, water content and impurity levels are seen over these time periods over this time period.

32. The data provided supports the stability of the novel food for a period of at least 9 months.

2.5 Specification

33. The specification parameters reported in Table 6 were assessed using internationally recognised methods or determined using internally developed and validated methods. The results of the analysis are detailed in Table 2 and indicate the novel food can be produced consistently to the specification.

Table 6: Specification of the novel food

Description
Cannabidiol is a white to slightly beige powder produced by chemical synthesis

Parameter	Specification	Method
Appearance	White to slightly beige crystalline powder	Visual inspection
Identity	Complies	IR: USP 197 HPLC: in-house method
Parameter	Specification	Method

Water content (%)	≤0.5	USP 921
Residue on ignition (%)	≤0.2	USP 281
Specific optical rotation (degrees)	-140 to -122	USP 781
CBD (%)	97 to 102	HPLC: in house method
Olivetol content (%)	≤0.15	HPLC: in house method
4-monobromo-cannabidiol (%)	≤0.15	HPLC: in house method
Delta-9-tetrahydrocannabidiol (THC) (%)	≤0.10	HPLC: in house method
Individual unspecified impurities (%)	≤0.15	HPLC: in house method
Total impurities (%)	≤1.0	HPLC: in house method
Residual solvents (mg/kg CBD)	-----	-----
Methanol	≤3000	GC: in house method
<i>n</i> -Heptane	≤5000	GC: in house method
Dichloromethane	≤600	GC: in house method
Triethylamine	≤5000	GC: in house method
Isopropanol	≤5000	GC: in house method
Isooctane	≤5000	GC: in house method

CBD = cannabidiol; GC = gas chromatography; HPLC = high-performance liquid chromatography; IR = infrared spectroscopy; USP = United States Pharmacopeia

34. Following discussion with the applicant it was agreed that information on microbiological and heavy metal limits did not need to be included in the specification to appropriately characterise the novel food. No specific recommendations were made to amend the specification. The ACNFP advised the information provided is sufficient to meet the specification of CBD and

appropriately characterises the novel food seeking authorisation.

2.6 History of Use

35. Hemp has been widely consumed in the UK and EU as a seed oil, in tea and as an alternative to hops in beer. Extracts of hemp including CBD and synthetic CBD have not been widely consumed and are considered novel foods. While CBD products are widely available on the UK high street, indicating some consumption of CBD as a food, at the time of publication, no previous applications for CBD have yet received authorisation as a novel food.
36. As detailed in the COT review of the literature there has been use of both hemp derived and synthetic forms of CBD for medicinal purposes. These provide an indication of the toxicological effects that should be explored in the testing regime – primarily effects on liver, thyroid and potential impacts on reproductive organs. Also reported are behavioural effects such as somnolence (sleepiness).
37. As reported in the COT review of the publicly available data on CBD and summary data on a medicinal product, signs of adverse effects on the liver were observed at doses of CBD as low as 5 mg/kg bw/day in patients and healthy human volunteers; this dose is equivalent to 350 mg in a 70 kg adult. The data in the literature also suggested that humans might be more sensitive to the adverse effects of CBD in the liver than laboratory animals.
38. Somnolence effects were noted at doses ≤ 10 mg/kg bw/day in human studies. Inhibitory drug-drug interactions have also been observed with some medications when CBD is co-administered at doses of 1 mg/kg bw/day (equivalent to 70 mg in a 70 kg adult); the likelihood of effects at lower doses has not been determined.
39. It is noted that the doses of CBD used for medicinal purposes are higher than those proposed for food use. The purpose of an assessment for medicines authorisation is different to that for food and this is reflected in the data requirements. Unlike medicines, there is no risk-benefit context in foods with

the requirement instead being that the products are safe. This means that outcomes that are considered to be an adverse event for food might be weighed differently in the context of the clinical benefits in a medicinal study.

40. Within the literature, further human studies using chemically derived CBD provides further evidence of a history of synthetic CBD use (Izegelov et al., 2010; Stereo Biotech Ltd., 2020; Klotz et al., 2019; Wheless et al., 2019). A review by Heuestis et al., 2019 of Cannabidiol Adverse effects and Toxicity notes that, while CBD is not risk-free, severe adverse events occur at doses higher than those recommended for human pharmacotherapies which are prescribed to treat forms of epilepsy. The data on previous consumption of CBD suggest areas for careful consideration in the toxicological review to understand potential effects at the lower doses used in foods.

2.7 Proposed Use and Anticipated Intake

41. The proposed use for the novel food is as a food supplement in the form of an oil sold as capsules and drops at the dose of 10 mg per day of CBD. These are detailed in Table 7. The intended use is food supplements as defined by UK legal requirements (The Food Supplements (England) Regulations 2003) as capsules, liquid or drops in dose form. A provisional acceptable daily intake (ADI) for the use of >98% pure form CBD has been established at 10 mg per day (October 2023) and discussed in the Toxicological information section.

Table 7: The proposed uses and maximum use levels for the novel food.

Food category	Maximum use level per day (mg CBD/day)
Food Supplements (for adults) as defined in the Food Supplements (England) Regulations 2003 as capsules, liquids or drops intended for those 18 years of age or over.	10

42. It is noted that there are already many products available containing CBD. As such, the assessment has been made on the basis of identification of a

maximum level of CBD that can be consumed per day per product. As such proposed uses will only be considered safe for all consumers within the assessment when below an intake maximum of 10mg of CBD per day.

43. Risk managers must consider whether consumers would benefit from information on the CBD content of foods in order to ensure their consumption does not exceed the provisional ADI of 10 mg per day for a healthy adult.
44. As recommended in the ACNFP and COT statement on CBD of > 98% purity, “The provisional ADI is recommended, subject to the existing advice to consumers that pregnant and breastfeeding women and people taking any prescription medication should avoid the consumption of CBD. Consumers on regular medications should seek advice from a medical professional before using any type of CBD food product. In addition, children and prospective parents trying for a baby are advised against consumption of CBD, as are those who are immunosuppressed, due to remaining data gaps and residual uncertainties concerning the safety of CBD for these groups of consumers.”
45. The food supplement products are to be labelled in accordance with the labelling requirements of Food Supplements (England) Regulations 2003as follows: Does not exceed the safe limit of 10 mg/day for a 70kg healthy adult. The novel food authorisation should also include the following: Not suitable for use under the age of 18. Not suitable for use during pregnancy or breastfeeding. If you are taking medication or have existing health conditions, please consult your doctor before using this product.
46. The ACNFP explored the potential for foreseeable misuse of the novel food. It is highlighted to risk managers that they may wish to consider whether risk management measures are needed beyond those in the food supplements regulation to ensure consumers are aware of the maximum safe dose for CBD intake per day and the excluded groups. The ACNFP also strongly recommended that risk managers consider how consumers can be supported to manage their intake appropriately within the safe limits identified and appreciate the nature of the potential risks at higher doses, for uses that are not in dosed forms.

2.8 Absorption, Distribution, Metabolism and Excretion (ADME)

47. The Absorption, Distribution, Metabolism and Excretion (ADME) of cannabidiol are known to be complicated by the food matrix in which the CBD is delivered and are currently still being defined by professional bodies.
48. The oral bioavailability of CBD is low, indicating that it is not absorbed to any notable extent following ingestion (Mechoulam et al., 2002). Published works report the bioavailability of CBD to be between 13 and 19% (Grotenhermen (2003)) or 6% (Hawsworth and McArdle (2004)). The low systemic availability was demonstrated by Martin-Santos et al., 2012 and further supported by a literature search which identified the pharmacokinetics of CBD in humans and concluded with dose dependent peak plasma concentrations of CBD and area under the curve results indicating minimal accumulation (Miller et al., 2018).
49. Following oral consumption, CBD is extensively metabolised in the liver. This rapid first pass metabolism contributes to the low oral bioavailability reported in the literature (Taylor et al., 2018; WHO, 2018). *In vitro* studies indicate that CYP3A4 and CYP2C19 are the primary hepatic enzymes responsible for first-pass metabolism of CBD; however, several other hepatic cytochrome P450 isoforms (CYP1A1, CYP1A2, CYP2C9, CYP2D6, and CYP3A5) have also demonstrated a capability of metabolising CBD (Jiang et al., 2011; Zendulka et al., 2016).
50. The metabolism of CBD is thought to follow two separate pathways. One is P450-mediated, in which CBD is metabolised into its major metabolite 7-COOH-CBD (which is a chemically inactive compound). This is followed by further metabolic reactions which yield the minor metabolites of CBD, including 6-OH-CBD (Devinsky et al., 2018; Taylor et al., 2018). The other involves decarboxylation (Kraemer et al., 2019). The resultant metabolites are predominantly excreted in faeces and urine (Hawsworth and McArdle, 2004; WHO, 2018).
51. Accumulation of CBD in plasma of up to 2-fold has been reported when steady state levels are compared with a single dose (such as in Taylor et al., 2018).

Additionally, minimal evidence of plasma accumulation has been reported in dosing studies over 5–9 days (Millar et al., 2018; Sellers et al., 2013; Stott et al., 2013).

52. The pharmacokinetics of CBD (24 studies) have also been systematically reviewed by Millar et al., (2018)., most of which assessed the administration of CBD at doses of 5–20 mg/day. This correlates to a low dose application similar to this CBD novel food application. With oral administration, single doses of 5.4 and 10 mg CBD achieved peak serum concentrations ($C_{\max}$) of 0.9 and 2.5 ng/ml. The time to maximum concentration ($T_{\max}$) was approximately 1 h, with a half-life between 1 and 3 hours. Given the intended use of this CBD, with an approximate half-life of one to three hours, with a total clearance of six hours, there are no significant concerns of accumulation.
53. Based on the information on ADME it is noted that, the bioavailability of CBD is typically low but can be affected by food matrix. This provides context for interpreting the toxicological data. It was noted that the potential for CBD to accumulate in the body has not been examined based on the data supplied. This also suggested the food context for the novel ingredient could impact whether the CBD present in the ingredient is more or less bioavailable. This has been taken into account in considering the assessment factors to account for uncertainty in setting the provisional ADI.

2.9 Nutritional information

54. The ACNFP sought clarification of the potential for the presence of antinutritional factors from the preparation. As the product is chemically derived from a well-defined synthetic process and characterisation indicated high CBD purity, no antinutritional factors were expected to be present. Information on the composition has confirmed that there is no presence of other components that would impact the digestion or absorption of nutrients from the diet.

55. The data on nutritional composition confirms that CBD has no caloric or nutritional value. The application is not intending that CBD replace another food in the diet. Consumption of the novel food at the proposed use levels is not expected to be nutritionally disadvantageous for consumers.

2.10 Toxicological information

56. Toxicological studies on CBD were performed by the applicant to support the safety assessment of the novel food. The respective study reports are unpublished and claimed as confidential and proprietary data. They were considered essential in the assessment of the safety of the novel food and were reviewed by the ACNFP. How data on systemic toxicity was managed and interpreted in the context of the provisional ADI is explained in the subchronic toxicology section below.

2.10.1 Genotoxicity

57. *In vitro* genotoxicity testing of CBD was conducted under Good Laboratory Practice (GLP) conditions and used the following OECD guidelines: *in vitro* bacterial reverse mutation test (OECD TG 471) and *in vitro* mammalian cell micronucleus test (OECD TG 487). This approach is recommended by the UK Committee on Mutagenicity and is also the basis of guidance on the preparation and submission of an application for authorisation of a novel food in the context of Regulation (EU) 2015/2283.

58. The *in vitro* bacterial reverse mutation test [(Gijsbrechts, 2020 (unpublished))] demonstrated that this CBD ingredient was non-mutagenic in the absence and presence of metabolic activation. In addition, the *in vitro* mammalian cell micronucleus test [(de Jong, 2020 (unpublished))] demonstrated that cannabidiol was non-clastogenic and non-aneugenic in the absence and presence of metabolic activation.

59. The results from these *in vitro* studies support the conclusion that the novel food (>98% pure CBD) is not genotoxic. This is consistent with the view of the Committee on Mutagenicity in reviewing CBD generically from evidence

available in the public domain (Committee on Mutagenicity; MUT/MIN/2020/1, 2020).

2.10.2 Sub-chronic toxicology study

60. The joint subgroup of the ACNFP and COT was formed to address a series of questions in relation to the safety of CBD, cannabinoids and hemp-derived ingredients.
61. A weight of evidence approach has allowed the Subgroup to identify a provisional ADI for CBD ingredients of >98% purity of 0.15 mg/kg bw/day or 10 mg per day for a 70 kg healthy adult (Joint position paper from the ACNFP and COT; FSA consumer advice published in October 2023). This value was identified to be protective of the most sensitive known effects in the liver and thyroid parameters and included consideration of data gaps and uncertainties.

2.10.3 Sub-chronic data supplied in this application

62. This applicant provided a Repeated Dose 90-Day Oral Toxicity Study in Rodents ([McGeoghie, 2020 unpublished]), which was conducted under GLP conditions and to OECD Technical Guideline 408. In this 90-day study, each group comprised 10 female and 10 male rats which were dosed with 0 (control – corn oil), 15, 36 or 72 mg/kg bw/day CBD once per day by oral gavage at a dose volume of 4 mL/ kg / body weight.
63. Review of the study supported the conclusion that it was of sufficient quality to support the safety of the novel food. The findings of the study were consistent with those considered in the development of the provisional ADI. It was, therefore, considered scientifically appropriate to apply the provisional ADI of 0.15 mg/kg bw/day or 10 mg/day as identified in the joint statement of the ACNFP and COT on >98% pure forms of CBD.

2.11 Allergenicity

64. This synthetic CBD comprises >98% pure CBD and the production process for CBD does not introduce any risk of allergenic potential. CBD as a chemical entity, suggests the potential for IgE mediated food allergy is unlikely.
65. The assessment considered whether the remaining 2% of the novel foods composition was likely to be allergenic or elicit food allergic reactions. It was noted that none of the raw materials or processing aids used in the production process are derived from or contain any of the allergenic food ingredients specified under Regulation (EU) No 1169/2011 of the European Parliament and of the Council of 25 October 2011 on the provision of food information to consumers. Suggesting the potential to elicit reactions in those sensitive to those foods is unlikely.
66. The novel food is unlikely to trigger allergic reactions in the target population under the proposed conditions of use.

3. Discussion

67. The novel food is a synthetic CBD ingredient containing $100\% \pm 2\%$ CBD (a Group A CBD ingredient), produced using a multi-step manufacturing process.
68. This synthetic CBD is intended to be used a food supplement for adults as an oil or as an oil in capsules at a dose of 10mg a day; it is not intended to replace any food.
69. In October 2023, the Joint ACNFP and COT subgroup identified a provisional acceptable daily intake (ADI) of 10 mg per day (0.15 mg/kg bw/day) for CBD novel foods containing 98% CBD or above, such as the novel food discussed in this assessment.
70. A weight of evidence approach was used to arrive at a provisional ADI of 10 mg/day (0.15 mg/kg bw/day). The most sensitive human health effects, that this provisional ADI protects against, are seen consistently in the liver and thyroid in

a number of studies using >98% pure CBD. This value also takes account of the lack of human-based long-term evidence and evidence regarding potentially vulnerable groups, which is applied here for this synthetic CBD⁵.

71. Based upon the dossier of evidence provided by the applicant, the safety of the novel food was reviewed and evidence to reach a conclusion on safety provided. The evidence presented is consistent with evidence presented to support the development of a provision ADI of 10 mg/day for CBD of 98% purity or above. As such the provisional ADI should be applied to this novel food.
72. This is subject to the existing advice to consumers that pregnant and breastfeeding women and people taking any prescription medication should avoid the consumption of CBD. Consumers on regular medications should seek advice from a medical professional before using any type of CBD food product. In addition, children and prospective parents trying for a baby are advised against consumption of CBD, as are those who are immunosuppressed, due to remaining data gaps and residual uncertainties concerning the safety of CBD for these groups of consumers. These contraindications would also apply to this novel food.
73. The maximum safe exposure for healthy adults of 70kg as identified in the provisional ADI is 10mg per day. If the inclusion level of this synthetic CBD leads to an intake per individual serving of each product type of 10 mg/day, only one product type per day should be consumed to ensure the provisional ADI is not exceeded. Multiple intakes of products containing CBD on the same day should be avoided to support minimising exposure to below the provisional ADI.

[5 Joint position paper from ACNFP & COT on establishing provisional ADI for pure form CBD in foods | Advisory Committee on Novel Foods and Processes](#)

4. Conclusion

The FSA and FSS have undertaken the assessment of synthetic CBD and both concluded that the novel food is safe under the proposed conditions of use and does not pose a safety risk to human health. The anticipated intake levels and the proposed use was not considered to be nutritionally disadvantageous.

These conclusions were supported by the information in the novel food dossier submitted by the applicant plus the supplementary information and could not have been reached without the following data claimed as proprietary by the applicant:

- annexes to the dossier which relate to the identity of the novel food, the production process, stability, methods of analysis, particle size analysis and toxicology.
- *in vitro* bacterial reverse mutation test [(Gijsbrechts, 2020 (unpublished))], *in vitro* mammalian cell micronucleus test [(de Jong, 2020 (unpublished))] and 90-day repeat dose feeding study [(McGeoghie, 2020 (unpublished))].

Acknowledgements

With thanks to the members of the ACNFP during the course of the assessment who were; Dr Camilla Alexander White, Dr Anton Alldrick, Dr Kimon Andreas Karatzas , Ms Alison Austin, Dr Mark Berry, Professor George Bassel, Dr Christine Bosch, Professor Dimitris Charalampopoulos, Dr Cathrina Edwards, Professor Susan Fairweather-Tait, Professor Paul Fraser, Dr Hamid Ghoddusi, Dr Andy Greenfield, Professor Wendy Harwood, Professor Huw D. Jones, Dr Ray Kemp, Dr Elizabeth Lund, Professor Harry J. McArdle, Rebecca McKenzie, Professor Clare Mills, Dr Antonio Peña-Fernández, Dr Lesley Stanley, Professor Hans Verhagen, Dr Maureen Wakefield, and Professor Bruce Whitelaw.

To note, interests were received from members of the ACNFP, Dr Alldrick declared a potential interest relating to his previous employment and this was considered a potential conflict and as a result he was not present for discussions of CBD by the

Committee. Emeritus Prof Harry McArdle declared an interest from his work with EFSA's novel food Committee in considering data requirements for CBD. While not seen as a conflict, to avoid Prof McArdle being subject to information that would influence his EFSA work, it was agreed that he would not be present in discussions for CBD by the ACNFP but could supply comments for consideration by the Committee upon review of the minutes.

References

Advisory Committee on Novel Foods and Processes (ACNFP) and Committee on Toxicity (COT), 2023. Joint position paper from the Advisory Committee on Novel Foods and Processes (ACNFP) & Committee on Toxicity (COT) on establishing a provisional acceptable daily intake (ADI) for pure form ($\geq 98\%$) cannabidiol (CBD) in foods, based on new evidence. Published October 2023 DOI: <https://doi.org/10.46756/sci.fsa.zcg392>

Committee On Mutagenicity Of Chemicals In Food, Consumer Products And The Environment, (MUT/MIN/2020/1), 2020; [Committee on Mutagenicity Meeting 2020/1](#)

de Jong BG, 2020 [unpublished]. Prepared by Charles River Laboratories Den Bosch B.V., DD 's Hertogenbosch, The Netherlands for Chanelle McCoy CBD Ltd., Loughrea, Galway, Ireland. Study Title: Cannabidiol: in vitro mammalian cell micronucleus test. Confidential. (Test Facility Study No. 20203946).

Devinsky O, Patel AD, Thiele EA, Wong MH, Appleton R, Harden CL, Greenwood S, Morrison G and Sommerville K, 2018. Randomized, dose-ranging safety trial of cannabidiol in Dravet syndrome [GWPCARE1 Part A Study Group]. *Neurology*, 90, e1204-e1211.

[EC, 2017. Commission Implementing Regulation \(EU\) 2017/2469 of 20 December 2017 laying down administrative and scientific requirements for applications referred to in Article 10 of Regulation \(EU\) 2015/2283 of the European Parliament and of the Council on novel foods.](#)

EFSA CONTAM Panel (Panel on Contaminants in the Food Chain), 2015. Scientific opinion on the risks for human health related to the presence of Tetrahydrocannabidiol (THC) in milk and other food from animal origin. DOI , [10.2903/j.efsa.2015.4141](https://doi.org/10.2903/j.efsa.2015.4141)

EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2021. Guidance on the preparation and submission of an application for authorisation of a

novel food in the context of Regulation (EU) 2015/2283 (Revision 1). Published 26 March 2021. <https://doi.org/10.2903/j.efsa.2011.2170>

[EU \(2015\) Regulation \(EU\) 2015/2283 of the European Parliament and of the Council of 25 November 2015 on novel foods, amending Regulation \(EU\) No 1169/2011 of the European Parliament and of the Council and repealing Regulation \(EC\) No 258/97 of the European Parliament and of the Council and Commission Regulation \(EC\) No 1852/2001](#)

Gijsbrechts JJA, 2020 [unpublished]. Prepared by Charles River Laboratories Den Bosch B.V., DD 's Hertogenbosch, The Netherlands for Chanelle McCoy CBD Ltd., Loughrea, Galway, Ireland. Study Title: Cannabidiol: bacterial reverse mutation test. Confidential. (Test Facility Study No. 20203944).

Grotenhermen, F. (2003) 'Pharmacokinetics and pharmacodynamics of cannabinoids', *Clinical Pharmacokinetics*, 42(4), 327–360.

<https://doi.org/10.2165/00003088-200342040-00003>

Hawksworth, G. and McArdle, K. (2004) 'Metabolism and pharmacokinetics of cannabinoids', in Guy, G, Whittle, B and Robson, P (eds.), *The Medicinal Uses of Cannabis and Cannabinoids*. London: London Pharmaceutical Press. pp. 205–228. Google Scholar

Huestis MA, Solimini R, Pichini S, Pacifici R, Carlier J, Busardò FP. Cannabidiol Adverse Effects and Toxicity. *Curr Neuropsychopharmacol*. 2019;17(10):974-989. doi:

[10.2174/1570159X17666190603171901](https://doi.org/10.2174/1570159X17666190603171901)

Jiang R, Yamaori S, Takeda S, Yamamoto I and Watanabe K, 2011. 'Identification of cytochrome P450 enzymes responsible for metabolism of cannabidiol by human liver microsomes', *Life Sciences*, 89(5–6), 165–170. doi:10.1016/j.lfs.2011.05.018.

Joint position paper from the Advisory Committee on Novel Foods and Processes (ANCFP) & Committee on Toxicity (COT) on establishing a provisional acceptable daily intake (ADI) for pure form (≥98%) cannabidiol (CBD) in foods, based on new evidence; October 2023. <https://doi.org/10.46756/sci.fsa.zcg392>

Izgelov D, Davidson E, Barasch D, Regev A, Domb AJ, Hoffman A. Pharmacokinetic investigation of synthetic cannabidiol oral formulations in healthy volunteers. Eur J Pharm Biopharm. 2020 Sep;154:108-115. doi: [10.1016/j.ejpb.2020.06.021](https://doi.org/10.1016/j.ejpb.2020.06.021)

Klotz KA, Grob D, Hirsch M, Metternich B, Schulze-Bonhage A, Jacobs J. Efficacy and Tolerance of Synthetic Cannabidiol for Treatment of Drug Resistant Epilepsy. Front Neurol. 2019 Dec 10;10:1313. doi: [10.3389/fneur.2019.01313](https://doi.org/10.3389/fneur.2019.01313)

Martin-Santos, R., Crippa, J. A., Batalla, A., Bhattacharyya, S., Atakan, Z., Borgwardt, S., Allen, P., Seal, M., Langohr, K., Farré, M., Zuardi, A. W and McGuire, P. K. (2012). Acute effects of a single, oral dose of d9-tetrahydrocannabinol (THC) and cannabidiol (CBD) administration in healthy volunteers. Current pharmaceutical design, 18(32), 4966–4979. <https://doi.org/10.2174/138161212802884780>

‘MDA’, Misuse of Drugs Act 1971, UK Public General Acts 1971 c. 38, [Misuse of Drugs Act 1971 \(legislation.gov.uk\)](https://www.legislation.gov.uk/ukpga/1971/38/contents/enacted)

Mechoulam R, Parker LA and Gallily R, 2002. Cannabidiol: an overview of some pharmacological aspects. Journal of Clinical Pharmacology, 42, 11S-19S.

Millar SA, Stone NL, Yates AS and O'Sullivan SE, 2018. A Systematic Review on the Pharmacokinetics of Cannabidiol in Humans. Frontiers in Pharmacology, 9, 1365 [13pp, plus supplementary table]. doi: [10.3389/fphar.2018.01365](https://doi.org/10.3389/fphar.2018.01365)

OECD (Organisation for Economic Co-operation and Development), 1997. Bacterial reverse mutation test. In OECD guidelines for the testing of chemicals. OECD guideline No 471 (updated & adopted: 21 July 1997). Paris, France: Organisation for Economic Co-operation and Development (OECD). <https://doi.org/10.1787/9789264071247-en>

OECD (Organisation for Economic Co-operation and Development), 1998. OECD principles of good laboratory practice. Series on principles of good laboratory practice and compliance monitoring, No. 1 (ENV/MC/CHEM(98) 17). Paris, France: Organisation for Economic Co-operation and Development (OECD), Environment Directorate, Chemicals Group and Management Committee.

<https://doi.org/10.1787/9789264078536-en>

OECD (Organisation for Economic Co-operation and Development), 2016. In vitro mammalian cell micronucleus test. In OECD guidelines for the testing of chemicals. OECD guideline No 487 (updated & adopted: 29 July 2016). Paris, France: Organisation for Economic Cooperation and Development (OECD).

<https://doi.org/10.1787/9789264264861-en>

-
OECD (Organisation for Economic Co-operation and Development), 2018. Repeated dose 90-day oral toxicity study in rodents. In OECD guidelines for the testing of chemicals. OECD guideline No 408 (updated and adopted 27 June 2018). Paris, France: Organisation for Economic Cooperation and Development (OECD).

<https://doi.org/10.1787/9789264070707-en>

McGeoghie S, 2020 [unpublished]. Prepared by Charles River Laboratories, East Lothian, UK for Chanelle McCoy CBD Ltd., Loughrea, Galway, Ireland. Study Title: Cannabidiol: a 90-day toxicity study by oral (gavage) in the rat. Confidential. (Test Facility Study No. 509549)

[Regulation \(EU\) No 1169/2011 of the European Parliament and of the Council of 25 October 2011 on the provision of food information to consumers, amending Regulations \(EC\) No 1924/2006 and \(EC\) No 1925/2006 of the European Parliament and of the Council, and repealing Commission Directive 87/250/EEC, Council Directive 90/496/EEC, Commission Directive 1999/10/EC, Directive 2000/13/EC of the European Parliament and of the Council, Commission Directives 2002/67/EC and 2008/5/EC and Commission Regulation \(EC\) No 608/2004 \(Text with EEA relevance\) \(legislation.gov.uk\)](#)

Sellers EM, Schoedel K, Bartlett C, Romach M, Russo EB, Stott CG, Wright S, White L, Duncombe P, Chen CF. A Multiple-Dose, Randomized, Double-Blind, Placebo-Controlled, Parallel-Group QT/QTc Study to Evaluate the Electrophysiologic Effects of THC/CBD Spray. Clin Pharmacol Drug Dev. 2013 Jul;2(3):285-94. doi: [10.1002/cpdd.36](https://doi.org/10.1002/cpdd.36)

Stott CG, White L, Wright S, Wilbraham D, Guy GW. A phase I study to assess the single and multiple dose pharmacokinetics of THC/CBD oromucosal spray. *Eur J Clin Pharmacol*. 2013 May;69(5):1135- 47. doi: [10.1007/s00228-012-1441-0](https://doi.org/10.1007/s00228-012-1441-0)

Stero Biotechs Ltd. Phase 2a Study to Evaluate the Safety, Tolerability and Efficacy of Cannabidiol as a Steroid-sparing Therapy in Steroid-dependent Crohn's Disease Patients. ClinicalTrials.gov Identifier: NCT04056442. Last Update Posted : June 30, 2020. Available online at: [A Phase 2a Study to Evaluate the Safety, Tolerability and Efficacy of Cannabidiol as a Steroid-sparing Therapy in Steroid-dependent Crohn's Disease Patients - Full Text View - ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT04056442)

[The Food Supplements \(England\) Regulations 2003, No. 1387, England](#)

[The Food Supplements \(Wales\) Regulations 2003 \(legislation.gov.uk\)](#)

[The Food Information Regulations 2014, No.1855](#)

Taylor L, Gidal B, Blakey G, Tayo B, Morrison G. A Phase I, Randomized, Double-Blind, Placebo Controlled, Single Ascending Dose, Multiple Dose, and Food Effect Trial of the Safety, Tolerability and Pharmacokinetics of Highly Purified Cannabidiol in Healthy Subjects. *CNS Drugs*. 2018 Nov;32(11):1053-1067. doi: [10.1007/s40263-018-0578-5](https://doi.org/10.1007/s40263-018-0578-5)

Wheless JW, Dlugos D, Miller I, Oh DA, Parikh N, Phillips S, Renfroe JB, Roberts CM, Saeed I, Sparagana SP, Yu J, Cilio MR; INS011-14-029 Study Investigators. Pharmacokinetics and Tolerability of Multiple Doses of Pharmaceutical-Grade Synthetic Cannabidiol in Pediatric Patients with Treatment-Resistant Epilepsy. *CNS Drugs*. 2019 Jun;33(6):593-604. doi: [10.1007/s40263-019-00624-4](https://doi.org/10.1007/s40263-019-00624-4)

[WHO \(World Health Organization\), 2018. Cannabidiol \(CBD\) critical review report. 40th Meeting - Expert Committee on Drug Dependence, Jun. 4-7, 2018, Geneva.](#)

Zendulka, O., Dovrtělová, G., Nosková, K., Turjap, M., Šulcová, A., Hanuš, L and Juřica, J. (2016). Cannabinoids and Cytochrome P450 Interactions. *Current drug metabolism*, 17(3), 206–226. <https://doi.org/10.2174/1389200217666151210142051>

Crown copyright 2023

This publication (not including logos) is licensed under the terms of the Open Government Licence v3.0 except where otherwise stated. Where we have identified any third party copyright information you will need to obtain permission from the copyright holders concerned.

For more information and to view this licence:

- visit [the National Archives website](#)
- email psi@nationalarchives.gov.uk
- write to: Information Policy Team, The National Archives, Kew, London, TW9 4DU

For enquiries about this publication, [contact the Food Standards Agency](#).



Follow us on Twitter: [@foodgov](#)



Find us on Facebook: facebook.com/FoodStandardsAgency

